

The Casimir energy of a massive fermion field revisited

F C Santos[†] and A C Tort^{‡,*}

Instituto de Física - Universidade Federal do Rio de Janeiro
C.P. 68528, Rio de Janeiro CEP 21945-970 Brasil

1st February 2008

Abstract

We introduce a general, simple, and effective method of evaluating the zero-point energy of a quantum field under the influence of arbitrary boundary conditions imposed on the field on flat surfaces perpendicular to a chosen spatial direction. As an example we apply the method to the Casimir effect associated with a massive fermion field on which MIT bag model type of boundary conditions are imposed.

Key words: Vacuum, zero-point oscillations, renormalization, Casimir effect
PACS: 11.10 -z

[†] e-mail: filadelf@if.ufrj.br

[‡] e-mail: tort@if.ufrj.br

* Present address: Institut d'Estudis Espacials de Catalunya (IEEC/CSIC) Edifici Nexus 201, Gran Capità 2-4, 08034 Barcelona, Spain. E-mail address: visit11@ieec.fcr.es

1 Introduction

The macroscopically observable vacuum energy shift associated with a quantum field is the regularised difference between the vacuum expectation value of the corresponding hamiltonian with and without the external conditions demanded by the particular physical situation at hand. At the one-loop level, when the external conditions are represented by boundary conditions this leads to the usual Casimir effect [1] – see Ref.[2] for an updated review on the theoretical and experimental aspects of this remarkable effect.

In the context of the Casimir effect, some configurations which depend on the type of quantum field, type of spacetime manifold and its dimensionality, specific boundary condition imposed on the quantum field on certain surfaces, lead to relatively simple spectra, others lead to more complex ones. The heart of the matter here is the evaluation of the spectral sum that results from definition of the Casimir energy. This evaluation requires regularization and renormalization, and recipes for accomplishing this range from the relatively simple cutoff method employed by Casimir himself [1] to the powerful and elegant generalised zeta function techniques [4]. Here, in the spirit of the representation of a spectral sum as a contour integral [5], we introduce a simple and effective way of evaluating the one-loop vacuum energy under external conditions based on well-known theorems of complex analysis, namely, the Cauchy integral formula and the Mittag-Leffler expansion theorem in one of its simplest versions. The method we present here is of sufficient generality so as to be successfully applied to a variety of cases. In this letter, however, we will limit ourselves to the example of a massive fermionic field under MIT boundary conditions and flat surface geometry, see Ref.[6] and references therein. The massless case was first calculated by Johnson [6] and as far as the present authors are aware of there is only one evaluation of this example of a massive fermionic Casimir energy due to Mamaev and Trunov [7]. This evaluation is only briefly sketched by these authors, therefore it seems interesting to test the method by providing an alternative derivation of this result.

2 The unregularised Casimir energy and a simple sum formula

Consider a quantum field in a $3+1$ dimensional flat spacetime under boundary conditions constraining the motion along one of the spatial directions, say, the $\mathcal{OX}_3$ -axis. At the one loop-level the (unregularised) Casimir energy will be given by

$$\mathcal{E} = \alpha \frac{\hbar c L^2}{2} \int \sum_n \frac{d^2 p_\perp}{(2\pi)^2} \Omega_n, \quad (1)$$

where α is a dimensionless factor that depends on the internal degrees of freedom of the quantum field under consideration, $p_\perp = \sqrt{p_1^2 + p_2^2}$, and

$$\Omega_n := \sqrt{p_\perp^2 + \frac{\lambda_n^2}{\ell^2} + m^2}, \quad (2)$$

where λ is the n -th real root of the transcendental equation determined by the boundary conditions, ℓ is a characteristic length along the $\mathcal{O}\mathcal{X}_3$ direction, and m is the mass of an excitation of the quantum field. A simple integral representation of $\sum_n \Omega_n$ can be written if we make use of Cauchy integral formula. In fact, it is easily seen that

$$\sum_n \Omega_n = - \oint_\Gamma \frac{dq}{2\pi} \sum_n \frac{2q^2}{q^2 + \Omega_n^2}, \quad (3)$$

where in principle Γ is a Jordan curve on the q -complex plane with $\Im q > 0$ consisting in a semicircle of infinitely large radius whose diameter is the entire real axis. Taking (3) into (1) we obtain

$$\mathcal{E} = -\alpha \frac{\hbar c L^2}{2} \int \frac{d^2 p_\perp}{(2\pi)^2} \oint_\Gamma \frac{dq}{2\pi} \sum_n \frac{2q^2}{q^2 + \Omega_n^2}. \quad (4)$$

In order to proceed we must be able to perform (at least formally) the discrete sum in (4).

Consider a complex function $G(z)$ of a single complex variable z , symmetrical on the real axis such that its roots are simple, non-zero and symmetrical with respect to the origin of the complex plane. The assumption that the origin is not a root of $G(z)$ is not a restrictive one because if $z = 0$ happens to be a root of $G(z)$ we can always divide $G(z)$ by some convenient power of z in order to eliminate zero from the set of the roots without introducing a new singularity. Let us order and count the roots of $G(z)$ in such a way that

$$\lambda_n = -\lambda_{-n}, \quad n \in Z - \{0\}. \quad (5)$$

Now define the following function

$$J(z) := \sum_{n \in Z - \{0\}} \frac{1}{z - i\lambda_n}. \quad (6)$$

The following properties of $J(z)$ are self-evident: (i) $J(z)$ has first order poles which are determined by the roots of $G(iz)$: (ii) the residua are all equal to unity. Taking into account property (5) we see that $J(z)$ can be rewritten in the form

$$\begin{aligned} J(z) &= \frac{1}{2} \left(\sum_{n \in Z - \{0\}} \frac{1}{z - i\lambda_n} + \sum_{n \in Z - \{0\}} \frac{1}{z + i\lambda_n} \right) \\ &= \sum_{n \in N} \frac{2z}{z^2 + \lambda_n^2}, \end{aligned} \quad (7)$$

where Z denotes the set of integers and N the set of the natural integers. Let us consider now the function $K(z) := G(iz)$ and state the following equality

$$J(z) = \frac{K'(z)}{K(z)}, \quad (8)$$

where the prime stands for the derivative with respect to z . In fact, the function $K'(z)/K(z)$ has the same simple poles as the originally defined $J(z)$ and the same residua, hence we can invoke the Mittag-Leffler theorem and state that (8) is true. It follows then that we can write

$$\frac{K'(z)}{K(z)} = \frac{d}{dz} \log K(z) = \sum_{n \in N} \frac{2z}{z^2 + \lambda_n^2}. \quad (9)$$

In order to make use of (9) we first link the dimensionless complex variable z to the auxiliary complex momentum variable q through the relation

$$q^2 + \Omega_n^2 = \frac{z^2 + \lambda_n^2}{\ell^2} \quad (10)$$

hence

$$z = \ell \sqrt{q^2 + p_\perp^2 + m^2}, \quad (11)$$

and we can write

$$\sum_n \frac{2q^2}{q^2 + \Omega_n^2} = \frac{\ell q^2}{z} \sum_n \frac{2z}{z^2 + \lambda_n^2}. \quad (12)$$

Upon changing variables ($d/dz = (z/\ell q)d/dq$), we obtain for the unregularised Casimir energy the following expression

$$\mathcal{E} = -\alpha \frac{\hbar c L^2}{2} \int \frac{d^2 p_\perp}{(2\pi)^2} \oint_\Gamma \frac{dq}{2\pi} q \frac{d}{dq} \log K(z), \quad (13)$$

which can be integrated by parts to yield

$$\begin{aligned} \mathcal{E} &= -\alpha \frac{\hbar c L^2}{2} \int \frac{d^2 p_\perp}{(2\pi)^2} \int_\Gamma \frac{dq}{2\pi} \frac{d}{dq} [q \log K(z)] \\ &+ \alpha \frac{\hbar c L^2}{2} \int \frac{d^2 p_\perp}{(2\pi)^2} \int_\Gamma \frac{dq}{2\pi} \log K(z). \end{aligned} \quad (14)$$

Notice that the integration is now performed on an open curve which lies on the Riemann surface of the integrand the projection of which on the q -complex plane is the curve Γ . The first term on the R.H.S of (14) contributes with a phase which cancels out with a phase coming from the second term since the final result is real. Hence, if we keep this in

mind the first term can be put aside and the unregularised Casimir energy will be given by

$$\mathcal{E} = \alpha \frac{\hbar c L^2}{2} \int \frac{d^2 p_\perp}{(2\pi)^2} \int_\Gamma \frac{dq}{2\pi} \log K(z). \quad (15)$$

Equation (15) is our main result. It is easily seen that it holds for arbitrary boundary conditions imposed on the field on flat surfaces perpendicular to the $\mathcal{OX}_3$ axis. It also holds when the $\mathcal{OX}_3$ direction is compactified by the imposition of topological conditions, periodic or antiperiodic, on the field along that direction.

3 The massive fermion field under MIT boundary conditions

As an example of the usefulness of (15) let us apply it to a massive fermion field under MIT boundary conditions imposed on field on the surface of an hypothetical bag consisting of two parallel square membranes perpendicular to the $\mathcal{OX}_3$ axis whose side a is much larger than ℓ , the distance between them. It can be shown that for this bag model the eigenvalues of p_3 are determined by the roots of the function $F(z \equiv p_3 \ell)$ defined by [7]

$$F(p_3 \ell) = \mu \sin(p_3 \ell) + p_3 \ell \cos(p_3 \ell), \quad (16)$$

where $\mu := m\ell$. Therefore we can choose $G(z)$ as

$$G(z) = \mu \frac{\sin z}{z} + \cos z \quad (17)$$

where we have divided $F(z)$ by z because $z = 0$ is a root of $F(z)$. We can easily prove that the roots of $G(z)$ are all real. Now we construct $K(z)$ according to

$$K(z) = G(iz) = \mu \frac{\sinh z}{z} + \cosh z. \quad (18)$$

Taking into account that for a fermionic quantum field in $3 + 1$ dimensions $\alpha = -4$, the vacuum energy is

$$\begin{aligned} \mathcal{E} &= -2\hbar c L^2 \int \frac{d^2 p_\perp}{(2\pi)^2} \int_\Gamma dq \ln \left[\frac{1}{2} \left(\frac{\mu}{z} + 1 \right) e^z \right] \\ &\quad - 2\hbar c L^2 \int \frac{d^2 p_\perp}{(2\pi)^2} \int_\Gamma dq \ln \left[1 + \frac{z - \mu}{z + \mu} e^z \right], \end{aligned} \quad (19)$$

where z is defined by (11). It is clear that the first term on the R.H.S of (19) is divergent, however, we can easily see also that this term is spurious in the sense that it carries a

contribution that does not depend on ℓ , is proportional to L^2 and represents the self-energy of the bag, plus another contribution proportional to the volume $L^2\ell$ of the bag representing a constant energy density present even if the bag were not there. Hence, we will subtract this term from (19) and write the regularised contribution as

$$\begin{aligned}\mathcal{E}^{(\text{reg})} &= -2\hbar c L^2 \int \frac{d^3p}{(2\pi)^3} \ln \left[1 + \frac{z - \mu}{z + \mu} e^{-2z} \right] \\ &= -\frac{\hbar c L^2}{\pi^2} \int_0^\infty dp p^2 \ln \left[1 + \frac{z - \mu}{z + \mu} e^{-2z} \right].\end{aligned}\quad (20)$$

Upon changing variables again the regularised Casimir energy can be rewritten as

$$\mathcal{E}^{(\text{reg})} = -\frac{\hbar c L^2}{\pi^2 \ell^3} \int_\mu^\infty dz z \sqrt{z^2 - \mu^2} \ln \left[1 + \frac{z - \mu}{z + \mu} e^{-2z} \right], \quad (21)$$

where now we have written $z^2 = \ell^2(p^2 + m^2)$ with $p^2 = q^2 + p_\perp^2$. This result is in agreement with the result obtained in [7]. For the case of massless fermions we obtain

$$\begin{aligned}\mathcal{E}^{(\text{reg})} &= -\frac{\hbar c L^2}{\pi^2 \ell^3} \int_0^\infty dz z^3 \ln [1 + e^{-2z}] \\ &= -\frac{7\pi^2 \hbar c L^2}{2880 \ell^3},\end{aligned}\quad (22)$$

which is the result obtained in [6]. Starting from (22) it is a straightforward matter to obtain the limits of the Casimir energy in the limits $\mu \rightarrow 0$ and $\mu \gg 1$, see [7].

4 Conclusions

In this letter we have derived a general expression, equation (15), for the evaluation of the Casimir energy of a quantum field in a flat manifold under the influence of boundary conditions imposed on the field on flat surfaces or topological conditions constraining the motion along a particular spatial direction. The present authors verified explicitly that equation (15) works well in several instances, in particular, for Robin boundary conditions results agree with those obtained by Romeo and Saharian [8]. This method can be extended and applied to cylindrical and spherical geometries embedded in $d + 1$ dimensional spacetimes. Here we have chosen the massive fermion field under MIT boundary conditions as an example and were able to rederive its associated vacuum energy in a clear cut way. Presently, this approach is being extended to more complex Casimir type problems.

Acknowledgments

One of the authors (A.C. Tort) wishes to acknowledge E. Elizalde and the hospitality of the Institut d'Estudis Espacials de Catalunya (IEEC/CSIC) and Universitat de Barcelona, Departament d'Estructura i Constituents de la Matèria where this work was completed, and the financial support of CAPES, the Brazilian agency for faculty improvement, Grant BExt 0168-01/2.

References

- [1] H.B.G. Casimir, Proc. K. Ned. Akad. Wet. **51**, (1948) 793.
- [2] M. Bordag, U. Mohideen and V.M. Mostepanenko: *New Developments in the Casimir effect*, quant-ph/0106045.
- [3] T.H. Boyer, Phys. Rev. **174**, (1968) 1764.
- [4] E. Elizalde, S.D. Odintsov, A. Romeo, A.A. Bytsenko and S. Zerbini, *Zeta function regularization techniques with applications*, (World Scientific, Singapore, 1994).
- [5] M. Bordag, E. Elizalde and K. Kirsten, J. of Math. Phys. 37, (1996) 895; M. Bordag, E. Elizalde, B. Geyer and K. Kirsten, Comm. Math. Phys. 179, (1996) 215; M. Bordag, E. Elizalde, K. Kirsten and S. Leseduarte, Phys. Rev. D 56, (1997) 4896; M. Bordag, E. Elizalde and K. Kirsten, J. of Phys. A 31, (1998) 1743.
- [6] K. Johnson, Acta Phys. Polonica **B6** (1975), 865.
- [7] S.G. Mamaev and N.N. Trunov, Izv. Vyssh. Uchebn. Zaved., Fiz. No 7, 9 (1980). English transl. Sov. Phys. **23** (1980) 551.
- [8] A. Romeo and A.A. Saharian, *Casimir effect for scalar fields under Robin boundary conditions on plates*, arXiv:hep-th/0007242 v1.